

RELATIVE TWISTING IN OUTER SPACE

MATT CLAY AND ALEXANDRA PETTET

ABSTRACT. Subsurface projection is indispensable to studying the geometry of the mapping class group and the curve complex of a surface. When the subsurface is an annulus, this projection is sometimes called *relative twisting*. We give two alternate versions of relative twisting for the outer automorphism group of a free group. We use this to describe sufficient conditions for when a folding path enters the *thin* part of Culler-Vogtmann's Outer space. As an application of our condition, we produce a sequence of fully irreducible outer automorphisms whose axes in Outer space travel through graphs with arbitrarily short cycles; we also describe the asymptotic behavior of their translation lengths.

1. INTRODUCTION

Culler and Vogtmann gave the first account of *Outer space* CV_k in their 1986 paper [20]: elements are finite marked projectivized metric graphs with fundamental group F_k , the rank k non-abelian free group, and two graphs are close when the lengths of some finite collection of elements of F_k are close. By considering the universal covers of the marked graphs, CV_k is also described as the space of free simplicial minimal isometric actions of F_k on \mathbb{R} -trees. Topologically, CV_k has the structure of a contractible simplicial complex (missing some faces) on which $\text{Out } F_k$ acts properly and simplicially by changing markings. Metrically, however, Outer space remains largely a mystery. Much of the conjectural picture for Outer space geometry comes from Teichmüller theory, where the Teichmüller metric, the Weil-Petersson metric, and the Thurston metric have been defined and extensively studied. Unfortunately Outer space lacks much of the structure that paves the way for these metrics; perhaps most notably, CV_k is *not* a manifold.

Of the three metrics on Teichmüller space mentioned above, only the third, the Thurston metric, has been interpreted in the Outer space setting; there it is more commonly referred to as the *Lipschitz metric*.

The first author is partially supported by NSF grant DMS-1006898. The second author was partially supported by NSF grant DMS-0856143, NSF RTG DMS-0602191, EPSRC grant EP/D073626/2, and an NSERC Discovery Grant.

Features of this metric were recorded by Francaviglia-Martino in [22, 23].

Algom-Kfir (see also Hamenstädt [26]) proved that axes of fully irreducible elements of $\text{Out } F_k$ are *strongly contracting*, so that CV_k exhibits a characteristic of negative curvature in these directions. Her result was anticipated by a theorem of Minsky [36], which showed that Teichmüller geodesics contained in the ϵ -*thick* part of Teichmüller space are strongly contracting, uniformly depending on ϵ . Algom-Kfir’s contraction constants depend on the outer automorphisms to which they belong. The question of whether these constants only depend on the geometry of the graphs along the axes has not been addressed.

For $\epsilon > 0$, we define CV_k^ϵ as the subset of CV_k consisting of graphs that contain a cycle of length less than ϵ . We should perhaps resist calling CV_k^ϵ the “thin part” of Outer space as it is not clear that Algom-Kfir’s theorem extends uniformly to geodesics in the complement of CV_k^ϵ . Nevertheless, this set does hold some nice properties analogous to those of the thin part of Teichmüller space; for instance, the cusps of the quotient $CV_k/\text{Out } F_k$ are contained in CV_k^ϵ , and the quotient $(CV_k - CV_k^\epsilon)/\text{Out } F_k$ of the complement is quasi-isometric to $\text{Out } F_k$.

The main results of this paper provide conditions, akin to those of Rafi [37] in the setting of Teichmüller space, that guarantee that a geodesic or an axis of a fully irreducible element travel through CV_k^ϵ . Our criteria are based on a notion of *relative twisting* in Outer space. We come at this from two different points of view, each motivated by the quest to find satisfactory analogues of subsurface projection and relative twisting from the theory of mapping class groups [35, 21].

Geometric: Our first approach to relative twisting directly adapts the original geometric definition to free groups. We give a pairing $\tau_a(G, G')$ between two graphs $G, G' \in CV_k$ relative to some nontrivial $a \in F_k$, which we define by means of the *Guirardel core* [24]. This is a certain 2-complex associated to the graphs that provides a means of selecting a geometry for F_k that “sees” both G and G' . We obtain a condition on the graphs that, when satisfied, enables us to construct a connecting geodesic between them, traveling through CV_k^ϵ .

Theorem 5.2. *Suppose $G, G' \in CV_k$ with $d = d_L(G, G')$ such that $\tau_a(G, G') \geq n+2$ for some $a \in F_k$. Then there is a geodesic $\alpha: [0, d] \rightarrow CV_k$ such that $\alpha(0) = G$ and $\alpha(d) = G'$ and for some $t \in [0, d]$, we have $\ell_{\alpha(t)}(a) \leq 1/n$. In other words, $\alpha([0, d]) \cap CV_k^{1/n} \neq \emptyset$.*

As a corollary, we get the following lower bound the distance between two marked graphs in CV_k .

Corollary 1.1. *Suppose $G, G' \in CV_k$ and G' does not have a cycle of length less than ϵ . Then:*

$$d_L(G, G') \geq \log \sup_{1 \neq a \in F_k} \epsilon \tau_a(G, G')$$

Proof. Let $a \in F_k$ be nontrivial. If $\tau_a(G, G') \geq n$, then by Theorem 5.2, there is a geodesic $\alpha: [0, d] \rightarrow CV_k$ such that $\alpha(0) = G$ and $\alpha(d) = G'$, and for some $t \in [0, d]$ that $\ell_{\alpha(t)}(a) \leq 1/n$. As G' does not have a cycle of length less than ϵ , it will follow from Proposition 2.1 that $d_L(\alpha(t), G') \geq \frac{\epsilon}{1/n} = \epsilon n$. As $\alpha(t)$ is on a geodesic from G to G' , the corollary holds. \square

The similar lower bound for Teichmüller space is a special case of a theorem of Rafi [38].

Algebraic: The second point of view to relative twisting gives a pairing $\tau_a(T, \Lambda)$ between a tree $T \in \overline{CV}_k$ and an algebraic lamination Λ of F_k relative to some nontrivial $a \in F_k$. This pairing measures how the axes of a in T overlap with the leaves of the lamination. It is similar to the notion of “twisting” used by Alibegović [3]. We obtain a criterion that implies that the axis of a fully irreducible element travels through CV_k^ϵ in terms of its unstable tree and lamination.

Theorem 5.3. *Suppose $\phi \in \text{Out } F_k$ is fully irreducible, with unstable tree T_- and lamination Λ_- such that $\tau_a(T_-, \Lambda_-) \geq n + 4$ for some $a \in F_k$. Then given any train-track G , there is an axis \mathcal{L}_ϕ for ϕ that contains G and a graph G_0 with $\ell_{G_0}(a) \leq 1/n$. In other words, $\mathcal{L}_\phi \cap CV_k^{1/n} \neq \emptyset$.*

As an application of Theorem 5.3, we examine outer automorphisms of F_k that are products of powers of two Dehn twists δ_1 and δ_2 which “fill” in an appropriate sense. We show (Section 6) that axes for $\delta_1^n \delta_2^{-n}$ travel through graphs with a cycle of length $\sim 1/n$. Moreover, we can estimate their translation lengths on CV_k ; we compute that they grow logarithmically in n (Theorem 6.6).

The proofs of Theorems 5.2 and 5.3 are similar. In both cases we show that large relative twist implies the existence of a certain path that contains a large power of a (Propositions 3.5 and 4.8). These paths, called vanishing paths, are folded, either in the map $G \rightarrow G'$ or in a train-track map representing ϕ , and are homotopically trivial in the image. The most efficient way to fold over a loop representing a several times is to first make a a short loop (Proposition 5.1).

It appears likely that our definition of algebraic twist (at least as used in Theorem 5.3) is a special case of our definition of geometric twist. We anticipate investigating this relationship in a further paper.

The paper is organized as follows. In Section 2 we review some of the basic theory of Outer space and the Lipschitz metric, irreducible outer automorphisms, train-track maps, and laminations; only Section 2.5 contains some new material. As this section is already lengthy, some background, such as a summary of currents for free groups, is suppressed until it is needed in Section 6. In Section 3, following an outline of some properties of Guirardel’s core and a brief review of relative twisting for the mapping class group, the first, “geometric,” analogue of relative twisting for $\text{Out } F_k$ is given. Section 4 is concerned with the second, “algebraic,” notion of relative twisting. Each of Sections 3 and 4 conclude with a proposition essential to the proofs of the main theorems, found in Section 5. In Section 6, we bring together results from Section 5 and previous papers of the authors [12, 13] to describe a method for constructing geodesic axes of fully irreducible elements which enter the thin part of Outer space.

Acknowledgments The authors would like to thank Kasra Rafi and Juan Souto for helpful conversations concerning this project. Additionally, the authors thank the referee for a careful reading of this work with helpful suggestions.

2. PRELIMINARIES

2.1. Outer space. We begin by fixing a generating set of the free group $F_k = \langle x_1, \dots, x_k \rangle$. Let G be a simplicial graph, i.e., a one-dimensional cell complex, with $\pi_1(G)$ isomorphic to F_k . Let R be a wedge of k (oriented) circles, with each circle identified to one of the generators of F_k . Then by a *marking* of G we mean a homotopy equivalence $\rho: R \rightarrow G$. From the map $\rho_*: \pi_1(R) \rightarrow \pi_1(G)$, we then have an identification of F_k with $\pi_1(G)$. Given two marked metric graphs $\rho_1: R \rightarrow G_1$ and $\rho_2: R \rightarrow G_2$, a map $f: G_1 \rightarrow G_2$ is a *change of marking* if it is linear on edges, and if $f \circ \rho_1: R \rightarrow G_2$ is homotopic to $\rho_2: R \rightarrow G_2$. A *topological representative* of $\phi \in \text{Out } F_k$ is a marked graph $\rho: R \rightarrow G$, together with a self homotopy equivalence $g: G \rightarrow G$, so that the homotopy equivalence $\rho^{-1} \circ g \circ \rho: R \rightarrow R$ induces ϕ on $\pi_1(R) = F_k$.

We denote by cv_k the *unprojectivized (Culler–Vogtmann) Outer space* consisting of marked metric graphs G , where $\pi_1(G) = F_k$ and the degree of every vertex of G is at least 3. Two points $\rho_1: R \rightarrow G_1$ and $\rho_2: R \rightarrow G_2$ in cv_k are equivalent if there is an isometry $\iota: G_1 \rightarrow G_2$ so that $\iota \circ \rho_1$ is homotopic to ρ_2 . An alternate description of cv_k is as the space of free minimal isometric F_k -actions on simplicial trees, and we will alternate freely between treating Outer space as a space of

trees and as a space of graphs. There is a right action of $\text{Out } F_k$ given by precomposing the marking (or F_k -action) by a representative of the outer automorphism. Outer space is defined as the projectivization of cv_k : $CV_k = cv_k/\mathbb{R}_{>0}$; it can be identified with the subspace of cv_k consisting of marked graphs whose edge lengths sum to 1.

To simplify the notation for elements in Outer space, we denote a marked metric graph $\rho: R \rightarrow G$ simply by G . A *path* in G is a continuous map $\alpha: I \rightarrow G$, where I is an interval of \mathbb{R} . For convenience, and when it is clear from context, α may denote either the map or its image in G ; while $[\alpha]$ will denote the image of α after ‘‘pulling it tight,’’ i.e., the image of any immersed homotopy (relative endpoints) representative of α . We then use $L_G(\alpha)$ to denote the length of $[\alpha]$ in G . For an element $a \in F_k$, we write $\ell_G(a)$ to denote the minimal length of a loop in G representing the conjugacy class of a .

For points $x, y \in T$, we use $[x, y]$ to denote the image of the unique tight path connecting x and y in T . For an F_k -tree T and an element $a \in F_k$, we write $\ell_T(a)$ to denote the minimal translation length of a in T . If $\ell_T(a) \neq 0$, then a has an invariant axis $T^{(a)}$ and $\text{vol}(T^{(a)}/\langle a \rangle) = \ell_T(a)$. If G is a graph with fundamental group F_k , then \tilde{G} denotes the universal cover of G , with a chosen base point so that there is an F_k -action on \tilde{G} . Clearly $\ell_G = \ell_{\tilde{G}}$.

Using the description as a space of tree actions, cv_k is topologized via the *axes topology*. That is, a tree $T \in cv_k$ is identified with a point in \mathbb{R}^{F_k} by the coordinates $(\ell_T(g))_{g \in F_k}$ [19]. Cohen and Lustig proved that the space of very small actions on \mathbb{R} -trees contains the closure \overline{cv}_k of cv_k [15]. The converse, that every very small minimal action on an \mathbb{R} -tree is the limit of free minimal simplicial actions, was shown by Bestvina and Feighn [6]. Recall that an action of F_k on an \mathbb{R} -tree is *very small* if arc stabilizers are trivial or maximal cyclic, and the stabilizer of any tripod is trivial.

Given two points G_1 and G_2 in the projectivized Outer space CV_k , let $f: G_1 \rightarrow G_2$ be a change of marking, and denote by $\sigma(f)$ the maximal slope of f (recall that f is linear on edges). We have the following proposition, due to White (see [1, 5]):

Proposition 2.1 (White). *Let G_1, G_2 be two graphs in CV_k . Then:*

$$\inf\{\sigma(f) \mid f: G_1 \rightarrow G_2 \text{ change of marking}\} = \sup_{1 \neq a \in F_k} \frac{\ell_{\tilde{G}_2}(a)}{\ell_{\tilde{G}_1}(a)}$$

Moreover both inf and sup are realized.

For G_1 and G_2 in $\text{Out } F_k$, let $\sigma(G_1, G_2)$ be the value in Proposition 2.1. We define a function $d_L : CV_k \times CV_k \rightarrow \mathbb{R}_{\geq 0}$ by

$$d_L(G_1, G_2) = \log \sigma(G_1, G_2).$$

Its only failure to be a distance is that it is not symmetric; it is not hard to construct examples of $G_1, G_2 \in CV_k$ with $d_L(G_1, G_2) \neq d_L(G_2, G_1)$ (see [2]). In spite of this anomaly, we will refer to d_L as the *Lipschitz metric* on CV_k . We remark that it is known that the minimal Lipschitz constant, taken over all continuous maps $f: G_1 \rightarrow G_2$ such that $f \circ \rho_1$ is homotopic to ρ_2 , is achieved by a map that is linear on edges ([1, 23]).

Example 2.2. We present an example of computing distances in CV_k that will be relevant to those examples constructed in Section 6. Fix a basis $\mathcal{T} = \mathcal{A} \cup \{t\}$ of F_k and an element $c \in \langle \mathcal{A} \rangle$ that is cyclically reduced with respect to \mathcal{T} . Consider the Cayley tree T and the marked graph $G = T/F_k$, metrized so that all edge lengths are equal to $1/k$.

Let δ be the automorphism that sends t to ct and acts as the identity on $\langle \mathcal{A} \rangle$. Then there is a change of marking map $f: G \rightarrow G\delta^n$ defined by subdividing the edge corresponding to t into $n+1$ edges and sending each of the first n edges over the edge path for c and the last edge over the edge corresponding to t . Therefore, the image of the edge t has length:

$$n\ell_G(c) + \frac{1}{k} = \frac{nk\ell_G(c) + 1}{k}$$

and hence the edge t has been stretched by $nk\ell_G(c) + 1$. Since the edge corresponding to t is the only edge stretched and since it is mapped to a tight loop we have that:

$$d_L(G, G\delta^n) = \log(nk\ell_G(c) + 1).$$

In the terminology from the proof of Theorem 2.1 in [1], the loop t is the subgraph $G_f \subset G$ and it is a legal loop.

The automorphism δ is an example of a *Dehn twist automorphism* (see Section 6); in this case corresponding to the Bass–Serre tree arising from the HNN-extension $\langle \mathcal{A}, c_0, t \mid t^{-1}ct = c_0 \rangle$. We refer to such a tree as a *cyclic tree*. For the case of a cyclic tree dual to an amalgamated free product $\langle \mathcal{A} \rangle *_{\langle c \rangle} \langle c, \mathcal{B} \rangle$ and its associated Dehn twist ($a \mapsto a, b \mapsto cbc^{-1}$), one can also show, using similar methods, that the distance from G to $G\delta^n$ is approximately $\log n$. In this case though, the obvious map sending edges corresponding to elements $b \in \mathcal{B}$ to $c^nb c^{-n}$ is not the optimal map. Instead one sends b to $c^{n/2}bc^{-n/2}$ and edges corresponding to elements $a \in \mathcal{A}$ to $c^{-n/2}ac^{n/2}$.

We remark for use in Section 6, that $d_L(G\delta^n, G) = d_L(G, G\delta^n)$. Indeed, $d_L(G\delta^n, G) = d_L(G, G\delta^{-n})$ and the same argument as above

shows that $d_L(G, G\delta^{-n}) = \log(nk\ell_G(c^{-1}) + 1)$. But of course $\ell_G(c) = \ell_G(c^{-1})$.

2.2. Bounded backtracking. Suppose that $f: T \rightarrow T'$ is a continuous map, where T and T' are trees. We say that f has *bounded backtracking* if there is a constant C such that for any path $[x, y] \subset T$ from x to y in T , and any $z \in [x, y]$, necessarily $d_{T'}([f(x), f(y)], f(z)) \leq C$. We denote by $BBT(f)$ the minimal such constant C . We note that for any given $T \in cv_k$ and $T' \in \overline{cv}_k$, that any F_k -equivariant map $f: T \rightarrow T'$ has bounded backtracking. Moreover $BBT(f) \leq \text{Lip}(f) \text{vol}_T(T/F_k)$, where $\text{Lip}(f)$ is the Lipschitz constant of the map f [7]. In particular, if T and T' are contained in the projectivized space CV_k , then $BBT(f) \leq \text{Lip}(f)$.

For a path $\alpha \subset T$, denote by $\alpha \dagger_L$ the path obtained by deleting the extremal paths of length L . The following is an easy consequence of bounded backtracking.

Lemma 2.3. *Suppose that $T, T' \in \overline{cv}_k$, that $f: T \rightarrow T'$ is an F_k -equivariant map that has bounded backtracking, and that $\ell: \mathbb{R} \rightarrow T$ is a parametrized geodesic. If $L > BBT(f)$, and if for some interval $I \subset \mathbb{R}$, a tight path $\alpha \subset T'$ is contained in $[f(\ell(I))] \dagger_L$, then necessarily $\alpha \subset [f(\ell(I'))]$ for any interval $I' \supset I$.*

Proof. Let $I = [x, y] \subset \mathbb{R}$ and assume that the hypotheses of the lemma hold, so that $d_{T'}(f(\ell(x)), \alpha) \geq L$ and $d_{T'}(f(\ell(y)), \alpha) \geq L$. Next let $x' \in I$ be such that $f(\ell(x'))$ is the endpoint of α closest to $f(\ell(x))$, and let $y' \in I$ be such that $f(\ell(y'))$ is the endpoint of α closest to $f(\ell(y))$.

Now suppose that $\alpha \not\subset [f(\ell(I'))]$ for some interval I' that contains I . As T' is a tree, either there exists an $x'' \in I' - I$ such that $f(\ell(x'')) = f(\ell(x'))$ or there exists an $y'' \in I' - I$ such that $f(\ell(y'')) = f(\ell(y'))$; without loss of generality, we assume the former. Then the path $[\ell(x'), \ell(x'')] \subset T$ and the point $\ell(x) \in [\ell(x'), \ell(x'')]$ violate bounded backtracking, as

$$d_{T'}([f(\ell(x')), f(\ell(x''))], f(\ell(x))) \geq L > BBT(f).$$

□

2.3. Irreducible elements and train-tracks. Let G be a graph. A *turn* is an unordered pair of oriented edges that share a common initial vertex. Letting \bar{e} denote the edge e with opposite orientation, we say that an edge path α *crosses* a turn $\{e_1, e_2\}$ if it contains an occurrence of either $\bar{e}_1 e_2$ or $\bar{e}_2 e_1$.

Let $g: G \rightarrow G$ be a homotopy equivalence that is linear on edges, mapping edges to edge paths. Then g induces a map on the set of turns

of G , as follows. Let v be the common initial vertex of the edges of a turn $\{e_1, e_2\}$. Some initial segment of e_1 is mapped onto an edge e'_1 based at $g(v)$, while some initial segment of e_2 onto an edge e'_2 , also based at $g(v)$. We assign $g(\{e_1, e_2\}) = \{e'_1, e'_2\}$.

The homotopy equivalence $g: G \rightarrow G$ is a *train-track map* if there is a collection \mathcal{LT} of turns such that:

- (1) \mathcal{LT} is closed under iteration of g and
- (2) for an edge $e \subset G$, any turn crossed by $g(e)$ is in \mathcal{LT} .

The unordered pairs of \mathcal{LT} are called *legal turns*, while an unordered pair of turns not in \mathcal{LT} is called an *illegal turn*. A path is *legal* if it only crosses legal turns. We will regularly refer to the underlying graph G as a “train-track.”

An element of $\phi \in \text{Out } F_k$ is *reducible* if some conjugacy class of a proper free factor of F_k is ϕ -periodic; otherwise ϕ is *irreducible*. Bestvina and Handel proved that every irreducible element of $\text{Out } F_k$ has a topological representative that is a train-track map [8]. If $g: G \rightarrow G$ is a train-track map representing an irreducible element of $\text{Out } F_k$, there is a metric on G such that g linearly expands each edge of G by the same factor λ , called the *expansion factor*. This factor is the Perron–Frobenius eigenvalue of the transition matrix for g ; a positive eigenvector for this eigenvalue specifies the metric on G .

All proper powers of a *fully irreducible* element ϕ of $\text{Out } F_k$ are irreducible. A fully irreducible element ϕ has the property that its minimal displacement in the Lipschitz metric is related to its expansion factor λ_ϕ by

$$\min_{G \in CV_k} d_L(G, G\phi) = \log(\lambda_\phi).$$

Moreover, this minimum is realized by a train-track map for ϕ . This relationship is one reason we choose not to symmetrize the Lipschitz metric, for typically the expansion factor of a fully irreducible element is not equal to that of its inverse. See for example [27].

For a fully irreducible element $\phi \in \text{Out } F_k$ and any tree $T \in CV_k$, the sequence $T\phi^n$ has a well-defined limit $T_+(\phi)$ in \overline{CV}_k , called the *stable tree* of ϕ [7, 33]. The *unstable tree* for ϕ , denoted by $T_-(\phi)$, is the stable tree for ϕ^{-1} , i.e., $T_-(\phi) = T_+(\phi^{-1})$. For an explicit description see [28]. We further note that if $T \in \overline{CV}_k - \{T_-(\phi)\}$, then $T\phi^n$ converges to $T_+(\phi)$ [33].

2.4. Geodesics in Outer space. Next, for an interval $I \subset \mathbb{R}$, we describe paths $I \rightarrow CV_k$ known as *folding lines*. We will be concerned with two types of such paths: those which connect two points G_1 and

G_2 in the interior CV_k , and those which are axes of fully irreducible elements. The latter were studied by Algom-Kfir in [1].

For $G_1, G_2 \in CV_k$, let $f: G_1 \rightarrow G_2$ be a change of marking map whose Lipschitz constant realizes $\sigma(G_1, G_2)$. Find a path $\tilde{\alpha}_1$ based at G_1 contained in an open simplex of (unprojectivized) cv_k , along which edges of G_1 shrink just until the map induced by f stretches every edge of the resulting graph by $\sigma(G_1, G_2)$; note that the lengths of those edges of G_1 that are stretched by exactly $\sigma(G_1, G_2)$ do not change along $\tilde{\alpha}_1$. Let the endpoint of the corresponding path α_1 in (projectivized) CV_k be H_1 , with the change of marking map $h: H_1 \rightarrow G_2$ induced by f . We choose a parameterization $\alpha_1: [0, d_L(G_1, H_1)] \rightarrow CV_k$ by arclength.

Now we construct a path $\alpha_2: [0, d_L(H_1, G_2)] \rightarrow CV_k$ with $\alpha_2(0) = H_1$ and $\alpha_2(d_L(H_1, G_2)) = G_2$. First subdivide the edges of H_1 to obtain a graph H'_1 so that the preimage of vertices in G_2 consists of vertices in H'_1 , while the induced map $h: H'_1 \rightarrow G_2$ remains cellular. Select a vertex v of H'_1 at which two edges e_1 and e_2 identified by h are based. Let $H(t)$ be the graph obtained from H'_1 by folding the initial segments of length $(1 - e^{-t})$ of e_1 and e_2 ; let $\alpha_2(t)$ be the graph in CV_k obtained from $H(t)$ by “forgetting” valence 2 vertices. Note that $d_L(H_1, \alpha_2(t)) = t$. Define $\alpha_2(t)$ in this way until e_1 and e_2 are completely identified; then repeat the above with the resulting graph. Continue this process until the map induced by f is an immersion in G_2 ; this is a finite process as H'_1 has a finite number of vertices. Note that the immersion is necessarily an isometry as every edge is stretched by the same factor; thus the fold line just constructed connects H_1 to G_2 in CV_k . Finally, let α be the concatenation of α_1 and α_2 ; a path based at G_1 and terminating at G_2 . Francaviglia and Martino [23, Theorem 5.5] proved that $\alpha: [0, d_L(G_1, G_2)] \rightarrow CV_k$ is a geodesic.

We can now describe a geodesic axis for a fully irreducible element ϕ of $\text{Out } F_k$. Let λ_ϕ be the expansion factor of ϕ , and let G be a train-track. To obtain a parametrized geodesic axis for ϕ , first find a folding path $\alpha: [0, \log(\lambda_\phi)] \rightarrow CV_k$ as above, connecting G to $G\phi$. Then define the graph $\alpha(t) = \alpha(t - n(t)\log(\lambda_\phi))\phi^{n(t)}$, where $n(t)$ is the integer $\lfloor \frac{t}{\log(\lambda_\phi)} \rfloor$, and let \mathcal{L}_ϕ denote the image of α . Algom-Kfir [1, Proposition 3.5] showed that $\alpha: \mathbb{R} \rightarrow CV_k$ is a geodesic parametrized by arclength.

2.5. Nielsen and vanishing paths. Suppose G is a graph with a homotopy equivalence $g: G \rightarrow G$. We make note of two special types of paths in G and collect some relevant results that will be useful for us in the sequel.

First, a path $\alpha \subset G$ is a *Nielsen path* if $[g(\alpha)] = [\alpha]$; it is *indivisible* if it is not a concatenation of nontrivial Nielsen paths, so that any Nielsen path is a concatenation of indivisible Nielsen paths.

Theorem 2.4 ([28], Corollary 2.14). *Suppose that $\phi \in \text{Out } F_k$ is fully irreducible with stable tree $T_+ = T_+(\phi)$, and that $g: G \rightarrow G$ is a train-track representative. Then there is a surjective F_k -equivariant map $f_g: \tilde{G} \rightarrow T_+$ such that if $[x, y] \subset \tilde{G}$ is a lift of a path $\alpha \subset G$ and $f_g(x) = f_g(y)$, then for some $m \geq 0$, the path $[g^m(\alpha)]$ is either a Nielsen path or trivial.*

Definition 2.5. Let $a \in F_k$ and $T \in \overline{cv}_k$ be such that $\ell_T(a) \neq 0$. We say that a path $\alpha \subset T$ (possibly infinite) *n-covers* a if $L_T(\alpha \cap T^{(a)}) \geq n\ell_T(a)$. In other words, α overlaps with a segment of the axis of a for length at least $n\ell_T(a)$; there is a point $x \in \alpha$ such that $a^n x \in \alpha$ as well.

Similarly, given $G \in CV_k$, we say a path $\alpha \subset G$ *n-covers* a if a lift of α to \tilde{G} *n-covers* a . In other words, α decomposes into $\alpha = \beta \cdot \alpha_0 \cdot \beta'$, where α_0 is the loop representing the conjugacy class of a^n .

Lemma 2.6. *Let ϕ and $g: G \rightarrow G$ be as in Theorem 2.4. Suppose that a Nielsen path α in G *n-covers* a , for some $a \in F_k$ with $\ell_{T_+}(a) > 0$ and $n \geq 2$. Then there is a subpath α_0 of α which *n-covers* a and which is contained in $\alpha' \dagger_\epsilon$ for some indivisible Nielsen path α' and $\epsilon > 0$.*

Proof. Suppose that α_0 is a shortest subpath of α that *n-covers* a , but is not contained in the interior of an indivisible Nielsen path.

We can express α as a concatenation $\alpha_1 \alpha_2 \dots \alpha_r$ of indivisible Nielsen paths α_i , $i = 1, \dots, r$. The g -fixed points of α are precisely the endpoints of the α_i 's. Then since α_0 is not contained in an indivisible Nielsen path, it must contain one of one of these fixed points p . It therefore contains at least n copies of the fixed point p . Therefore some sequence $\alpha_i \dots \alpha_j$ forms a Nielsen path and corresponds to the conjugacy class of a . A closed Nielsen path corresponds to a periodic loop. It follows that $\ell_{T_+}(a) = 0$, contradicting the hypothesis on a . \square

A path $\alpha \subset G$ is a *vanishing path* of g if $[g^m(\alpha)]$ is trivial (i.e., is a point) for some $m \geq 1$. We record the following observation from [4]:

Lemma 2.7. *Let ϕ and $g: G \rightarrow G$ be as in Theorem 2.4, and suppose that α is an indivisible Nielsen path. Then any subpath $\beta \subseteq \alpha \dagger_\epsilon$ for some $\epsilon > 0$ is contained in a vanishing path.*

Proof. An indivisible Nielsen path can be decomposed into a sequence of legal paths as $\alpha = \alpha_0 \cdot \beta_0 \cdot \bar{\beta}_1 \cdot \bar{\alpha}_1$, where $g(\alpha_i) = \alpha_i \cdot \beta_i$ and $g(\beta_0) = g(\beta_1)$ [8]. Hence for $\epsilon > 0$, with large enough n , the path $[g^n(\alpha \dagger_\epsilon)]$ is contained in $\beta_0 \cdot \bar{\beta}_1$, and hence $[g^{n+1}(\alpha \dagger_\epsilon)]$ is trivial. \square

Putting together Theorem 2.4 and Lemmas 2.6 and 2.7, we have the following:

Proposition 2.8. *Suppose that $\phi \in \text{Out } F_k$ is fully irreducible with stable tree $T_+ = T_+(\phi)$, that $g: G \rightarrow G$ is a train-track representative of ϕ , and that $a \in F_k$ is such that $\ell_{T_+}(a) > 0$. Then there is a surjective F_k -equivariant map $f_g: \tilde{G} \rightarrow T_+$ such that if $[x, y] \subset \tilde{G}$ is a lift of a reduced path $\alpha \subset G$ that n -covers a , for some $n \geq 2$, and if $f_g(x) = f_g(y)$, then α contains a vanishing path that n -covers a .*

2.6. Laminations and the map \mathcal{Q} . There are several notions of a ‘‘lamination’’ on a free group. For a full discussion on three different approaches and the relations between them, see [16, 17, 18]. We will only briefly describe the elements of the theory we need here.

The group F_k is hyperbolic and hence has a boundary ∂F_k . We denote:

$$\partial^2 F_k = \{(x_1, x_2) \in \partial F_k \times \partial F_k \mid x_1 \neq x_2\}$$

This set is naturally identified with the space of oriented *bi-infinite geodesics* in a tree $T \in cv_k$ as we explain now.

An oriented bi-infinite geodesic is an isometric embedding $\ell: \mathbb{R} \rightarrow T$ considered up to reparametrization preserving the orientation. Any geodesic has two distinct endpoints in ∂T , denoted $\ell(\infty)$ and $\ell(-\infty)$. We can thus identify the geodesic ℓ with endpoints $(\ell(\infty), \ell(-\infty)) \in \partial^2 T = \{(x_1, x_2) \in \partial T \times \partial T \mid x_1 \neq x_2\}$, which, via the action of F_k , is naturally identified with $\partial^2 F_k$. Conversely, a point $(x_1, x_2) \in \partial^2 F_k$ determines two distinct points $x'_1, x'_2 \in \partial T$. Between these two points, there is a unique (up to orientation preserving reparametrization) oriented geodesic $\ell: \mathbb{R} \rightarrow T$ such that $\ell(\infty) = x'_1$ and $\ell(-\infty) = x'_2$. We will use this identification without further remark.

There is fixed point free involution on $\partial^2 F_k$ defined by $\sigma: (x_1, x_2) \rightarrow (x_2, x_1)$, corresponding to reversing a geodesic’s orientation in T .

A *lamination* is a closed F_k -invariant and σ -invariant subset $\Lambda \subseteq \partial^2 F_k$. The set of algebraic laminations inherits a Hausdorff topology from $\partial^2 F_k$, which is described in [16]. A nontrivial element $a \in F_k$ determines a *minimal rational* lamination:

$$\Lambda(a) = \{(ga^{-\infty}, ga^{+\infty}) \cup (ga^{+\infty}, ga^{-\infty}) \mid g \in F_k\}$$

Note that the set $\Lambda(a)$ depends only on the conjugacy class of a . Although we will not need them here, we mention that the set of *rational* laminations consists of finite unions of minimal rational laminations. The most important example of a lamination in what follows is the *stable lamination* $\Lambda_+(\phi)$ associated to a fully irreducible element

$\phi \in \text{Out } F_k$, as defined in [7].¹ The *unstable lamination* $\Lambda_-(\phi)$ associated to ϕ is the stable lamination of ϕ^{-1} , so that $\Lambda_-(\phi) = \Lambda_+(\phi^{-1})$.

Let $g: G \rightarrow G$ be a train-track representative of ϕ with expansion factor λ . After passing to a power of g if necessary, we can assume that g has a fixed point x contained in the interior of an edge. For some small ϵ -neighborhood U of x , we have that $g(U) \supset U$. Fix an isometry $\ell: (-\epsilon, \epsilon) \rightarrow U$ and extend this to the unique isometric immersion $\ell: \mathbb{R} \rightarrow G$ such that $\ell(\lambda^n t) = g^n(t)$. This immersion lifts to a collection of geodesics $\tilde{\ell}: \mathbb{R} \rightarrow \tilde{G}$. Using the identification mentioned above between $\partial^2 F_k$ and the space of geodesics in \tilde{G} , the collection of all geodesics (called *leaves*) constructed as above determines a closed F_k -invariant subset of $\partial^2 F_k$ called the *stable lamination*. It is proved in [7] that this set is well-defined independent of g . The leaves of $\Lambda_+(\phi)$ are *quasi-periodic* [7], so that for every $L > 0$ there is an $L' > L$ such that for every interval I of length L and every interval I' of length L' there is an element $x \in F_k$ such that $x\ell(I) \subseteq \ell(I')$.

Given a basis \mathcal{A} of F_k and a tree $T \in \overline{cv}_k$, define the set $L_{\mathcal{A}}^1(T)$ as the set of right infinite reduced words $x = x_1 x_2 x_3 \cdots$ in the basis \mathcal{A} such that for some $p \in T$, the sequence of points $(x_1 x_2 \cdots x_i)p$ is bounded. The identification of right infinite reduced words in \mathcal{A} with ∂F_k identifies $L_{\mathcal{A}}^1(T)$ with a subset $L^1(T) \subseteq \partial F_k$ that is well-defined independent of the choice of basis. Bounded backtracking ensures the existence of a well-defined injective map $\mathcal{Q}: \partial F_k - L^1(T) \rightarrow \partial T$. Using the injectivity of \mathcal{Q} on $\partial F_k - L^1(T)$, we associate to any oriented bi-infinite geodesic $\alpha = (x_1, x_2) \in \partial^2 F_k - (L^1(T))^2$ an oriented bi-infinite geodesic $\alpha_T \subset T$, namely $\alpha_T = (\mathcal{Q}(x_1), \mathcal{Q}(x_2)) \in \partial^2 T$, if neither endpoint of α is in $L^1(T)$; otherwise we define α_T to be the empty set. In the latter case, following [33], we say that the geodesic α is *T-bounded*.

In certain cases, the map $\mathcal{Q}: \partial F_k - L^1(T) \rightarrow \partial T$ extends to a map on ∂F_k .

Proposition 2.9 ([33], Proposition 3.1). *Suppose $T \in \overline{cv}_k$ has dense orbits and trivial arc stabilizers (e.g., the stable tree for a fully irreducible outer automorphism). There exists a map $\mathcal{Q}: L^1(T) \rightarrow \overline{T}$ to the metric closure \overline{T} of T such that, for any $f: T_0 \rightarrow T$, where $T_0 \in cv_k$, and any ray ρ in T_0 representing $x \in L^1(T)$, the point $\mathcal{Q}(x)$ belongs to the closure of $f(\rho)$ in \overline{T} .*

¹Note that in [28], the stable lamination is called the “expanding lamination” and denoted by Λ_- as it is more naturally associated to $T_-(\phi)$. See Proposition 2.10.

Combining this with the previous discussion, we have a map $\mathcal{Q}: \partial F_k \rightarrow \bar{T} \cup \partial T$ whenever T has dense orbits and trivial arc stabilizers.

The relation between stable trees and laminations is illustrated by the following.

Proposition 2.10 ([7], Lemma 3.5 (3) & [33], Corollary 2.3). *Suppose that $\phi \in \text{Out } F_k$ is fully irreducible with stable tree T_+ and unstable lamination Λ_- . Let $\mathcal{Q}: \partial F_k \rightarrow \bar{T} \cup \partial T$ be the map defined following Proposition 2.9. Then for any leaf $\ell \in \Lambda_-$, we have $\mathcal{Q}(\ell(\infty)) = \mathcal{Q}(\ell(-\infty))$.*

In light of the above propositions, we can define for any tree $T \in \bar{cv}_k$ with dense orbits and trivial arc stabilizers [17]:

$$L_{\mathcal{Q}}^2(T) = \{(x_1, x_2) \in \partial^2 F_k \mid \mathcal{Q}(x_1) = \mathcal{Q}(x_2)\}$$

where $\mathcal{Q}: \partial F_k \rightarrow \bar{T} \cup \partial T$ is the map from Proposition 2.9. With this definition, Proposition 2.10 states that $\Lambda_-(\phi) \subseteq L_{\mathcal{Q}}^2(T_+^\phi)$ where $\phi \in \text{Out } F_k$ is fully irreducible. If ϕ is hyperbolic (i.e., ϕ does not have nontrivial periodic conjugacy class) then $L_{\mathcal{Q}}^2(T_+^\phi)$ is the “diagonal closure” of $\Lambda_-(\phi)$ [30].

Missing from the above is a discussion of *measured geodesic currents* and *Dehn twist automorphisms* needed for Section 6. We defer their discussion until needed.

3. GEOMETRIC RELATIVE TWISTING

Our first definition of relative twisting for $\text{Out } F_k$ follows closely the original geometric notion for the mapping class group, upon replacing a surface with a suitable 2-complex. This complex, the *Guirardel Core* for two F_k -trees T, T' , is a certain F_k -invariant subspace $\mathcal{C} \subset T \times T'$ (with the diagonal action). We will not need the precise definition of the complex for our purposes; rather, we record in Proposition 3.1 just those properties of \mathcal{C} we do require, together with references.

In the following, if p is a point in T , then $\mathcal{C}_p = \{x' \in T' \mid (p, x') \in \mathcal{C}\}$; similarly, for $p' \in T'$, we have $\mathcal{C}_{p'} = \{x \in T \mid (x, p') \in \mathcal{C}\}$. These sets are called the *slices of the core*.

Proposition 3.1. *Suppose $T, T' \in CV_k$.*

- (1) *The core $\mathcal{C} \subset T \times T'$ is nonempty, connected, closed, CAT(0), F_k -invariant and has convex fibers, i.e., the slices \mathcal{C}_p and $\mathcal{C}_{p'}$ are each convex for all $p \in T$ and $p' \in T'$. Moreover, \mathcal{C} is the minimal (with respect to inclusion) subset of $T \times T'$ with these properties [24, Main Theorem].*

- (2) The quotient \mathcal{C}/F_k has finite volume [24, Theorem 8.1]. This volume is called the intersection number, denoted $i(T, T')$.
- (3) For any $p' \in T'$ that is not a vertex, any arc $\gamma \subset \mathcal{C}_{p'}$ is contained in a vanishing path of any change of marking map $f: T \rightarrow T'$ [4, Lemma 3.7 & Remark 5.3].

For the complete definition of the core, along with examples, see [4, 24].

Before going further we briefly recall relative twisting for curves on a surface. Let S be a surface of genus at least two, equipped with a hyperbolic metric. We can consider $\pi_1(S)$ as a discrete group of isometries of \mathbb{H}^2 , so that $S = \mathbb{H}^2/\pi_1(S)$. Fix three simple closed curves, α, β, γ , so that β and γ both intersect α . Each of these curves corresponds to a conjugacy class of an element in $\pi_1(S)$, and we can assume that all three are geodesics on S . Let S_α be an annular cover of S corresponding to α ; that is, the quotient of \mathbb{H}^2 by the cyclic group generated by a representative of α in $\pi_1(S)$. We let α_c denote the unique lift of α to S_α that is closed. The *twist of β and γ relative to α* is defined as the maximum geometric intersection number between β' and γ' that intersect $\alpha_c \subset S_\alpha$, where β' and γ' range over lifts of β and γ to S_α .

We can reformulate this in terms of the universal cover \tilde{S} , defining the relative twist as follows. Fixing a lift $\tilde{\alpha}$ of α to \tilde{S} , the twist of β and γ relative to α is the maximum number of α -translates of $\tilde{\gamma}$ that intersect $\tilde{\beta}$, over all choices of lifts $\tilde{\beta}, \tilde{\gamma}$ of β, γ that intersect $\tilde{\alpha}$. See Figure 1. This interpretation can be extended to trees in Outer space, using the Guirardel core of F_k -trees T and T' in place of \tilde{S} .

The role of the simple closed curves β, γ is filled by tracks on \mathcal{C} , and of the simple closed curve α by the axis of an element of F_k in \mathcal{C} . A *track for T* in \mathcal{C} is the set $\{p\} \times \mathcal{C}_p$ where p is the midpoint of some edge of T ; a track for T' is defined similarly. We will also use *track* to refer to the image of a track in the quotient \mathcal{C}/F_k . We record some elementary properties of tracks.

- (1) Every track separates \mathcal{C} .
- (2) Every track is a finite subtree.
- (3) Every track is a convex subset of \mathcal{C} .

As \mathcal{C} is CAT(0), and every nontrivial element acts hyperbolically, the minset of a nontrivial element $g \in F_k$ is isometric to a product $Y \times \mathbb{R}$, where Y is a convex subset of \mathcal{C} [10]. An *axis* of a nontrivial element $a \in F_k$ is a subset of the minset of a of the form $\{y_0\} \times \mathbb{R}$. The element a acts by translation on any of its axes.

Lemmas 3.2 and 3.3 describe the extent to which the intersection number between tracks and axes is well-defined.




FIGURE 1. The relative twist of β and γ with respect to α is 5.

Lemma 3.2. *Let $a \in F_k$ be a nontrivial element, $T, T' \in CV_k$ and consider the core $\mathcal{C} \subset T \times T'$. If a track τ in \mathcal{C} intersects an axis of a in \mathcal{C} , then it intersects every axis of a in \mathcal{C} .*

Proof. Let $\tau = \{p\} \times \mathcal{C}_p$ be a track for T that intersects an axis for a . Let $Y \times \mathbb{R} \subset \mathcal{C}$ be the minset for a . As tracks and axes are convex, their intersection is convex as well, and hence connected. Moreover, as tracks are finite, there are $s, t \in \mathbb{R}$ such that $(Y \times \mathbb{R}) \cap \tau \subset Y \times [s, t]$.

Let $\{y_0\} \times \mathbb{R} \subset Y \times \mathbb{R}$ be an axis of a that intersects τ . As $(\{y_0\} \times \mathbb{R}) \cap \tau$ is connected, we have that $\{y_0\} \times (-\infty, s)$ and $\{y_0\} \times (t, \infty)$ project to different components of $T - \{p\}$. Hence τ separates $\{y_0\} \times \{s-1\}$ from $\{y_0\} \times \{t+1\}$ in \mathcal{C} . If there were an axis, say $\{y_1\} \times \mathbb{R}$, that did not intersect τ , then the concatenation of the fiber-wise paths

$$\{y_0\} \times \{s-1\} \rightarrow \{y_1\} \times \{s-1\} \rightarrow \{y_1\} \times \{t+1\} \rightarrow \{y_0\} \times \{t+1\}$$

would be a path that connected $\{y_0\} \times \{s-1\}$ to $\{y_0\} \times \{t+1\}$ avoiding τ , which is a contradiction as τ separates these points. Thus every axis for a intersects τ . \square

Lemma 3.3. *Let $a \in F_k$ be a nontrivial element, $T, T' \in CV_k$ and consider the core $\mathcal{C} \subset T \times T'$. Fix a track τ' for T' that intersects an*

axis of a in \mathcal{C} . Let τ_0 and τ_1 be two tracks for T that intersect an axis of a . Then:

$$\left| |\tau' \cap \langle a \rangle \tau_0| - |\tau' \cap \langle a \rangle \tau_1| \right| \leq 1.$$

Proof. Suppose $|\tau' \cap \langle a \rangle \tau_0| = n$. We will show that $|\tau' \cap \langle a \rangle \tau_1| \geq n - 1$. The statement of the lemma follows after interchanging τ_0 and τ_1 and applying the same argument.

Since a track for T and a track for T' can intersect at most once in \mathcal{C} , the track τ' intersects exactly n $\langle a \rangle$ -translates of τ_0 . We claim that there is an i such that τ' intersects $a^{i+j}\tau_0$ for $j = 0, \dots, n-1$. Indeed, this follows as τ' is connected, and as $a^m\tau_0$ separates $a^{m-r}\tau_0$ from $a^{m+s}\tau_0$ for all m and positive r and s . Replacing τ_0 by $a^i\tau_0$, we can assume that τ' intersects $\tau_0, \dots, a^{n-1}\tau_0$.

If $\tau_1 = a^i\tau_0$ for some i , then the statement is obvious. Otherwise, as $|\tau' \cap \langle a \rangle \tau_1|$ only depends on the orbit of τ_1 under a , we can replace τ_1 by $a^i\tau_1$ for some i to assume that τ_1 separates τ_0 from $a\tau_0$.

We claim that τ' intersects $\tau_1, \dots, a^{n-2}\tau_1$. Indeed, since τ' is connected and since $a^i\tau_1$ separates $a^i\tau_0$ from $a^{i+1}\tau_0$, both of which intersect τ' for $i = 0, \dots, n-2$, the track τ' must intersect $a^i\tau_1$. \square

As $|\tau' \cap \langle a \rangle \tau| = |\langle a \rangle \tau' \cap \tau|$, Lemma 3.3 shows that if τ'_0 and τ'_1 are tracks for T' that intersect an axis of a , and likewise τ_0 and τ_1 are tracks for T that intersect an axis of a , then:

$$\left| |\tau'_0 \cap \langle a \rangle \tau_0| - |\tau'_1 \cap \langle a \rangle \tau_1| \right| \leq 2.$$

With this bound we can define the relative twist number. By $Tr_a(T)$ we denote the set of tracks for T in \mathcal{C} that intersect an (and hence every) axis of a . We define the set $Tr_a(T')$ similarly.

Definition 3.4. Given $T, T' \in CV_k$ and a nontrivial element $a \in F_k$, define the *twist of T and T' relative to a* as:

$$\tau_a(T, T') = \max_{\tau' \in Tr_a(T'), \tau \in Tr_a(T)} |\tau' \cap \langle a \rangle \tau|.$$

We remark that this number is always finite. Indeed, as tracks are finite, the quantities we are maximizing over are finite. Then by the above discussion, there are only finitely many distinct possibilities for these numbers.

The significance of the relative twist number to the geodesic in CV_k connecting two marked graphs is the following, to be used in Section 5 to prove Theorem 5.2:

Proposition 3.5. *Suppose $G, G' \in CV_k$ with $d = d_L(G, G')$ such that $\tau_a(G, G') \geq n$ for some nontrivial $a \in F_k$. Then for every change of*

marking map $g: G \rightarrow G'$, there is a vanishing path $\gamma \subset G$ that n -covers a .

Proof. Let \tilde{G} and \tilde{G}' be the universal covers of G and G' respectively, and consider the core $\mathcal{C} \subset \tilde{G} \times \tilde{G}'$. Fix an axis of a , and tracks $\tau \in Tr_a(\tilde{G})$, $\tau' \in Tr_a(\tilde{G}')$ such that $\tau_a(G, G') = |\tau' \cap \langle a \rangle \tau|$.

Let m_0 and m_1 be the least and greatest integer, respectively, such that $\tau' \cap a^{m_0} \tau \neq \emptyset$ and $\tau' \cap a^{m_1} \tau \neq \emptyset$. Thus $\tau_a(G, G') = m_1 - m_0 \geq n$. Denote $x_0 = \tau' \cap a^{m_0} \tau$ and $x_1 = \tau' \cap a^{m_1} \tau$ and let ρ be the path in τ' connecting x_0 to x_1 . As $\tau' = \mathcal{C}_{p'} \times \{p'\}$ for some point $p' \in \tilde{G}'$, we can consider ρ as a path in $\mathcal{C}_{p'} \subset \tilde{G}$. Notice that the endpoints of ρ , also denoted x_0 and x_1 , are on the axis for a , and that $a^{m_1 - m_0} x_0 = x_1$. Thus a is n -covered by ρ . By Proposition 3.1(3), the path ρ is contained in a vanishing path γ for any change of marking map $G \rightarrow G'$. As γ contains ρ , the vanishing path γ n -covers a as well. \square

4. ALGEBRAIC RELATIVE TWISTING

In this section we give our algebraic interpretation of relative twisting and develop some consequences that will be useful for applications in later sections. The key result is Proposition 4.8, which is used to prove Theorem 5.3.

Definition 4.1. Given $T \in \overline{cv}_k$, a lamination $\Lambda \subset \partial^2 F_k$, and an element $a \in F_k$, if $\ell_T(a) \neq 0$, then we define the *twist of T and Λ relative to a* to be:

$$\tau_a(T, \Lambda) = \sup \left\{ \frac{L_T(\alpha_T \cap T^{(a)})}{\ell_T(a)} \mid \alpha \in \Lambda \right\}.$$

If $\ell_T(a) = 0$, then define $\tau_a(T, \Lambda) = 0$.

Recall that given $\alpha = (x_1, x_2) \in \partial^2 F_k$, we have that α_T is equal to $(\mathcal{Q}(x_1), \mathcal{Q}(x_2)) \in \partial^2 T$ if α is not T -bounded, and is equal to the empty set otherwise. We insist that $L_T(\emptyset) = 0$.

Remark 4.2. We allow for the possibility that $\tau_a(T, \Lambda) = \infty$. This occurs in particular for the rational lamination $\Lambda(a)$, when $\ell_T(a) \neq 0$.

Central to our analysis is the following proposition:

Proposition 4.3. *Suppose $a \in F_k$, $T \in \overline{cv}_k$, and that Λ is a lamination containing no T -bounded geodesic. If $\{T_i\}$ is a sequence of trees in cv_k converging to T , and $\{\Lambda_i\}$ is a sequence of laminations converging to Λ , then:*

$$\lim_{i \rightarrow \infty} \tau_a(T_i, \Lambda_i) \geq \tau_a(T, \Lambda).$$

Proof. The proposition is obviously true when $\ell_T(a) = 0$, and so we assume that $\ell_T(a) > 0$.

We proceed with the following:

Claim. *If $T \in \overline{cv}_k$, and if $\alpha, \beta \in \partial^2 F_k$ are not T -bounded and $\alpha(\infty), \alpha(-\infty), \beta(\infty), \beta(-\infty)$ are four distinct points, then for sufficiently close $T' \in cv_k$, we have $L_{T'}(\alpha_{T'} \cap \beta_{T'})$ close to $L_T(\alpha_T \cap \beta_T)$.*

Proof. As \mathcal{Q} is injective on $\partial F_k - L^1(T)$, we have that $\alpha_T \cap \beta_T$ is compact set.

Fix a tree $T_0 \in cv_k$, a map $f: T_0 \rightarrow T$ that is linear on edges, and elements $a_i^\pm \in T_0$ so that $[a_i^-, a_i^+] \rightarrow \alpha_{T_0}$. Then $\alpha_i = [f(a_i^-), f(a_i^+)] \rightarrow \alpha_T$; in particular, the overlap of α_i and α_T can be made arbitrarily large. Similarly define $b_i^\pm \in T_0$ and $\beta_i = [f(b_i^-), f(b_i^+)]$ so that, as before, we have $\beta_i \rightarrow \beta_T$.

Fix a $T' \in cv_k$ and an equivariant map $f': T_0 \rightarrow T'$, linear on edges. As before, we have $\alpha'_i = [f'(a_i^-), f'(a_i^+)] \rightarrow \alpha_{T'}$ and $\beta'_i = [f'(b_i^-), f'(b_i^+)] \rightarrow \beta_{T'}$.

Now choose n large enough so that each of $\alpha_n \cap \alpha_T$ and $\beta_n \cap \beta_T$ contains $\alpha_T \cap \beta_T$. Then increase n if necessary so that $\alpha'_n \cap \alpha_{T'}$ and $\beta'_n \cap \beta_{T'}$ each contain $\alpha_{T'} \cap \beta_{T'}$. For sufficiently close trees T, T' , the lengths $L_T(\alpha_n \cap \beta_n)$ and $L_{T'}(\alpha'_n \cap \beta'_n)$ of the overlaps are close [11, 25]. By choice of n , we have $L_T(\alpha_n \cap \beta_n) = L_T(\alpha_T \cap \beta_T)$ and $L_{T'}(\alpha'_n \cap \beta'_n) = L_{T'}(\alpha_{T'} \cap \beta_{T'})$. The claim follows. \square

We are now prepared to complete the proof of the proposition. First assume that $\tau_a(T, \Lambda) \neq \infty$. This implies that no geodesic in Λ has $a^{+\infty}$ or $a^{-\infty}$ as an endpoint and so we can use the above claim. Let ϵ be small and choose a geodesic $\alpha \in \Lambda$ so that:

$$L_T(\alpha_T \cap T^{(a)}) / \ell_T(a) > \tau_a(T, \Lambda) - \epsilon.$$

Then by the above:

$$\frac{L_{T'}(\alpha_{T'} \cap T'^{(a)})}{\ell_{T'}(a)} > \frac{L_T(\alpha_T \cap T^{(a)})}{\ell_T(a)} - \epsilon > \tau_a(T, \Lambda) - 2\epsilon$$

for T' sufficiently close to T . For Λ' sufficiently close to Λ , there exists $\alpha' \in \Lambda'$ so that $\alpha_{T'} \cap T'^{(a)} \subset \alpha'_{T'}$. Thus:

$$\tau_a(T', \Lambda') \geq \frac{L_{T'}(\alpha_{T'} \cap T'^{(a)})}{\ell_{T'}(a)} > \tau_a(T, \Lambda) - 2\epsilon$$

for T' sufficiently close to T and Λ' sufficiently close to Λ . Since we obtain such an inequality for every ϵ , the proposition holds.

Suppose $\tau_a(T, \Lambda) = \infty$. If $a^{+\infty}$ or $a^{-\infty}$ is an endpoint of a geodesic in Λ , then $\tau_a(T', \Lambda) = \infty$ for all trees $T' \in cv_k$. Else, we have that for, for every $M > 0$ there is a geodesic $\alpha \in \Lambda$ so that:

$$\infty > L_T(\alpha_T \cap T^{(a)})/\ell_T(a) > M.$$

Then arguing in a similar fashion as above, we have for T' sufficiently close to T and Λ' sufficiently close to Λ :

$$\tau_a(T', \Lambda') \geq \frac{L_{T'}(\alpha_{T'} \cap T'^{(a)})}{\ell_{T'}(a)} > \frac{M}{2}.$$

Since we obtain such an inequality for every M , the proposition holds. \square

Remark 4.4. Examples of tree, lamination pairs satisfying the hypotheses of Proposition 4.3 are:

- (1) $T_+(\phi), \Lambda_+(\phi)$ where $\phi \in \text{Out } F_k$ is fully irreducible,
- (2) T, Λ where $T \in cv_k$ and Λ is any lamination, and
- (3) $T, \Lambda(a)$ whenever $\ell_T(a) \neq 0$.

For a fully irreducible element ϕ with large twist $\tau_a(T_-(\phi), \Lambda_-(\phi))$ for some nontrivial $a \in F_k$, our goal is to locate a train-track G_0 of ϕ with a vanishing path that n -covers a , similar to Proposition 3.5. Our tool to produce such a path is Proposition 2.8. First we see how to use the lamination to get the required setup.

Lemma 4.5. *Suppose $\phi \in \text{Out } F_k$ is fully irreducible, $g: G \rightarrow G$ is a train-track representative for ϕ , $T_+ \in \overline{cv}_k$ is the stable tree for ϕ , $f_g: \tilde{G} \rightarrow T_+$ is the induced map from Theorem 2.4, and $\ell: \mathbb{R} \rightarrow \tilde{G}$ is a leaf of the unstable lamination Λ_- . Then for all $I \subset \mathbb{R}$, there exists $I' = [x, y] \subset \mathbb{R}$ such that $I \subseteq I'$ and $f_g(\ell(x)) = f_g(\ell(y))$.*

Proof. We claim that for any $L \geq 0$, there is an interval $[a, b] \subseteq \mathbb{R}$ such that $|b - a| \geq L$ and $f_g(\ell(a)) = f_g(\ell(b))$. The lemma follows: by the quasi-periodicity of ℓ , there is then an interval $I_0 = [a, b] \subset \mathbb{R}$ such that $f_g(\ell(a)) = f_g(\ell(b))$ and $x\ell(I) \subseteq \ell(I_0)$. Setting $I' = \ell^{-1}(x^{-1}\ell(I_0))$ completes the proof. We must then just establish the claim.

Since ℓ is a leaf of the unstable lamination, we have $\mathcal{Q}(\ell(-\infty)) = \mathcal{Q}(\ell(\infty))$. There are sequences $a_i \rightarrow -\infty$ and $b_i \rightarrow \infty$ such that $f_g(\ell(a_i)) \rightarrow \mathcal{Q}(\ell(-\infty)) = \mathcal{Q}(\ell(\infty))$ and $f_g(\ell(b_i)) \rightarrow \mathcal{Q}(\ell(\infty))$ [33, Lemma 3.4]. Now we have two cases, either the sequences $\{f_g(\ell(a_i))\}$ and $\{f_g(\ell(b_i))\}$ are in the same component of $\overline{T}_+ - \{\mathcal{Q}(\infty)\}$ or they are not.

If the sequences are in the same component, choose n with $|b_n - a_n| > L$. The arc α connecting $f_g(a_n)$ to $f_g(b_n)$ is then disjoint from $\mathcal{Q}(\ell(\infty))$,

and there is a unique point $p \in \alpha$ which is closest to $\mathcal{Q}(\ell(\infty))$. As \overline{T}_+ is an \mathbb{R} -tree, p is on the geodesic $[f_g(a_n), \mathcal{Q}(\ell(-\infty))]$. Then by continuity of $f\ell$, there is an $a' \leq a_n$ such that $f_g(a') = p$. Likewise, there is a $b' \geq b_n$ such that $f_g(b') = p$. Thus the interval $[a', b']$ satisfies the claim.

Now suppose the sequences are not in the same component. If $f_g(\ell(\mathbb{R}))$ crosses the point $\mathcal{Q}(\ell(\infty))$ infinitely many times, then we can find a sequence of points $a_i, b_i \in \mathbb{R}$ such that $f_g(\ell(a_i)) = f_g(\ell(b_i)) = \mathcal{Q}(\ell(\infty))$ such that $|b_i - a_i| \rightarrow \infty$. Indeed, there is a lower bound on the distance between two pre-images of $\mathcal{Q}(\infty)$ in \mathbb{R} as every edge of \tilde{G} is isometrically embedded by f_g . For large enough i , the interval $[a_i, b_i]$ satisfies the claim.

Finally, suppose $f_g(\ell(\mathbb{R}))$ crosses $\mathcal{Q}(\ell(\mathbb{R}))$ only finitely many times. Let a be the smallest number such that $f_g(\ell(a)) = \mathcal{Q}(\ell(\infty))$. Then arguing as in the first case using sequences $a_i \rightarrow -\infty$ and $b_i \rightarrow a$ ($b_i < a$) we can find the desired interval. \square

In the next proposition, we find a candidate vanishing path in a train-track G that folds over a several times. The technicalities in its proof arise from the fact that, as the hypothesis concerns the *unstable* lamination, we must first find a large power of a covered by a leaf of the lamination in a train-track map for the *inverse* ϕ^{-1} of ϕ . Care is then required in mapping this leaf over to a train-track for ϕ , as there might be excessive cancellation. We resolve this difficulty by applying powers of ϕ^{-1} , so that the length of a dominates any such cancellation.

Proposition 4.6. *Suppose that ϕ is fully irreducible with unstable tree T_- and lamination Λ_- , with $\tau_a(T_-, \Lambda_-) \geq n+2$ for some $a \in F_k$. Then there exists a train-track graph $G \in CV_k$ for ϕ and a leaf of the unstable lamination $\ell: \mathbb{R} \rightarrow \tilde{G}$ such that for all $L > 0$, there is a finite interval $I \subset \mathbb{R}$ such that $[\ell(I)]\dagger_L$ n -covers a .*

Proof. Let $H \in CV_k$ be a train-track graph for ϕ^{-1} with train track representative $h: H \rightarrow H$ and $G \in CV_k$ a train track graph for ϕ with train-track map $g: G \rightarrow G$. Fix Lipschitz homotopy equivalences $\kappa: H \rightarrow G$ and $\kappa': G \rightarrow H$ representing the change in markings. Thus the following diagram is commutative up to homotopy:

$$\begin{array}{ccc} G & \xrightarrow{g} & G \\ \kappa \uparrow & & \downarrow \kappa' \\ H & \xleftarrow{h} & H \end{array}$$

Notice that κ lifts to $\tilde{\kappa}: \tilde{H} \rightarrow \tilde{G}$, with bounded backtracking. In particular, we can pick a constant C such that if a path $\gamma \subset H$ has length at least C , then $\kappa(\gamma)$ is not homotopically trivial relative to its endpoints.

As $\tau_a(T_-, \Lambda_-) \geq n + 2$, Proposition 4.3 implies that $\tau_a(\tilde{H}_0, \Lambda_-) \geq n + 2$, where $H_0 = H\phi^{-M}$ for some large M . Define $G_0 = G\phi^{-M}$. Now there is a leaf $\ell: \mathbb{R} \rightarrow \tilde{H}_0$ of the unstable lamination and an interval $I_0 \subset \mathbb{R}$ such that $\ell(I_0) \subset \tilde{H}_0^{(a)}$ and $L_{\tilde{H}_0}(\ell(I_0)) \geq (n + 2)\ell_{\tilde{H}_0}(a)$. Notice that this implies that the loop representing the conjugacy class of a in H_0 is legal with respect to h . Then if we let λ be the expansion factor for ϕ^{-1} , and let $d = \ell_{\tilde{H}_0}(a)$, we have $\ell_{\tilde{H}_0\phi^{-m}}(a) = \lambda^m d$. Let N be such that $\lambda^N d \geq C$. Define $G_1 = G_0\phi^{-N}$ and $H_1 = H_0\phi^{-N}$.

Fix $L > 0$ and let L' be such that if we have paths $\gamma \subset \gamma'$ in H_0 and γ has length at least L' then the path $\kappa(\gamma)$ intersects $[\kappa(\gamma')]$ in a path of length at least L (necessarily a subpath of $[\kappa(\gamma')]$, but not necessarily a subpath of $[\kappa(\gamma)]$). Indeed, such an L' exists as κ is a quasi-isometry and the graph G_0 has valence bounded by $2k$. Finally, extend I_0 to $I = I_1 \cup I_0 \cup I_2 \subset \mathbb{R}$ by including intervals I_1 and I_2 of length at least $(L' + C)/\lambda^N$.

Notice that the lengths of $h^N(\ell(I_1))$ and $h^N(\ell(I_2))$ in H_1 are at least $L' + C$. Moreover, the initial subsegment $\iota \subset h^N(\ell(I_1))$ of length L' maps by κ to a segment in G_1 that intersects $[\kappa h^N \ell(I)]$ in a path of length at least L , so that $\kappa(\iota)$ does not cancel with $\kappa h^N \ell(I_0)$, as ι and $h^N \ell(I_0)$ are separated in $h^N \ell(I)$ by a subsegment of length at least C . Similarly for the terminal subsegment of $h^N(I_2)$.

Recall $\ell(I_0)$ $(n + 2)$ -covers a in H_0 . This clearly implies that $h^N \ell(I_0)$ $(n + 2)$ -covers a in H_1 as all of the paths are legal with respect to h . Thus $h^N \ell(I_0)$ contains a subpath whose image in H_1 represents the conjugacy class of a^{n+2} . As $\ell_{\tilde{H}_1}(a) \geq C$, a subsegment of length $n\ell_{\tilde{G}_1}(a)$ in $[\kappa h^N \ell(I_0)]$ survives after tightening $[\kappa h^N \ell(I_1)] \cdot [\kappa h^N \ell(I_0)] \cdot [\kappa h^N \ell(I_2)]$ to get $[\kappa h^N \ell(I)]$. Now tighten $\kappa \ell$ to get a leaf of the unstable lamination $\ell_1: \mathbb{R} \rightarrow \tilde{G}_1$. Thus $L_{\tilde{G}_1}([\ell_1(I)] \dagger_L \cap \tilde{G}_1^{(a)}) \geq n\ell_{\tilde{G}_1}(a)$ and hence $[\ell_1(I)] \dagger_L$ n -covers a . \square

Proposition 4.8 will now follow relatively quickly once we observe the following consequence of Proposition 4.6.

Corollary 4.7. *Suppose that ϕ is fully irreducible with unstable tree T_- and lamination Λ_- , such that $\tau_a(T_-, \Lambda_-) \geq n + 2$ for some $a \in F_k$. Then there exists a train-track $G \in CV_k$ for ϕ , with train-track map $g: G \rightarrow G$, and a path $\gamma = [x, y] \subset \tilde{G}$ such that:*

- (1) γ n -covers a ; and

- (2) $f_g(x) = f_g(y)$ where $f_g: \tilde{G} \rightarrow T_+$ is the induced map (see Theorem 2.4) from \tilde{G} to T_+ , the stable tree for ϕ .

Proof. Let G be the train-track graph given by Proposition 4.6, and let $f_g: \tilde{G} \rightarrow T_+$ be the induced map. Let $\ell: \mathbb{R} \rightarrow \tilde{G}$ be the leaf of the unstable lamination and $I \subset \mathbb{R}$ the interval produced by Proposition 4.6, for $L = BBT(f_g) + 1$. For this I , let I' be the interval given by Lemma 4.5.

We claim that $\gamma = [\ell(I')]$ satisfies the conclusion of the corollary. Indeed by Lemma 2.3, as $[\ell(I)]\dagger_L$ contains a path in the axis of a of length $n\ell_{\tilde{G}}(a)$, so does $\gamma = [\ell(I')]$. By construction, f_g identifies the endpoints of γ . \square

Proposition 2.8 applied to the train-track G and the path γ of Corollary 4.7 give the following:

Proposition 4.8. *Suppose ϕ is a fully irreducible element with unstable tree T_- and lamination Λ_- such that $\tau_a(T_-, \Lambda_-) \geq n + 2$ for some $a \in F_k$. Then there exists a train-track $G \in CV_k$ for ϕ , with train-track map $g: G \rightarrow G$, and a vanishing path $\gamma \subset G$ that n -covers a . \square*

5. FINDING SMALL CYCLES

With our definition of relative twist, we can prove the analogue of a special case of Rafi's characterization of short curves along geodesics in Teichmüller space [37].

Proposition 5.1. *Suppose $G, G' \in CV_k$, $f: G \rightarrow G'$ is a change of marking map with minimal slope, $d = d_L(G, G')$ and $a \in F_k$. If there is a vanishing path $\gamma \subset G$ that $(n + 2)$ -covers a , then there is a geodesic $\alpha: [0, d] \rightarrow CV_k$ such that $\alpha(0) = G$, $\alpha(d) = G'$ and for some $t \in [0, d]$, we have $\ell_{\alpha(t)}(a) \leq 1/n$.*

Proof. Shrinking the edges of G such that each edge is stretched by exactly e^d results in a marked graph G_0 and provides a geodesic $\alpha: [0, d_0] \rightarrow CV_k$ such that $\alpha(0) = G$, $\alpha(d_0) = G_0$ and $d = d_L(G_0, G') + d_0$. Denote the induced map $G_0 \rightarrow G'$ by f .

Consider the graph $H_a = \tilde{G}_0^{(a)}/\langle a \rangle$ and the map $h_{G'}: H_a \rightarrow G'$ which is the composition of the immersion $H_a \rightarrow G_0$ with the map $f: G_0 \rightarrow G'$. By appropriately subdividing and folding the graph H_a , after pruning we obtain a graph immersion $H'_a \rightarrow G'$. Now choose a folding path based at G_0 whose folds correspond to the folds performed on H_a . The end of the folding path is a graph G_1 in which H'_a is immersed.

Let $d_1 = d_L(G_0, G_1)$. As f has minimal slope, the above path extends α to a geodesic $\alpha: [0, d_1] \rightarrow CV_k$ such that $\alpha(0) = G$ and $\alpha(d_1) = G_1$ [23, Theorem 5.5]. Further $d_L(G_1, G') = d - d_1$. Denote the induced map $G_1 \rightarrow G'$ by f_1 .

The geodesic segment α can further be extended to a geodesic by folding G_1 . The image of γ in G_1 (which we denote by γ_1) is a vanishing path for the map f_1 .

Claim. *The path $\gamma_1 \subset G_1$ n -covers a .*

Proof of Claim. Consider the graph $\tilde{G}_0/\langle a \rangle$. This graph consists of a collection of trees attached to H_a . We consider the graph H_a as oriented counterclockwise and decompose H_a into subsegments $\delta_1 \epsilon_1 \cdots \delta_\ell \epsilon_\ell$ where the δ_i are the maximal subsegments that remain upon folding $H_a \rightarrow H'_a$ and pruning. The images of the ϵ_i are what get pruned. There is a lift of γ to $\tilde{G}_0/\langle a \rangle$ that decomposes into subpaths $\gamma = \beta_0 \alpha \beta_1$ where β_0 and β_1 are embedded and α is the immersed path that covers H_a $n + 2$ times.

When folding the segments ϵ_i , the initial part of β_1 may (by equivariance) become identified with some portion of H_a . We are interested in bounding how much is identified with the terminal portion of α as this could reduce the amount of γ_1 that covers a . We will show that the portion of α identified is a segment of H_a . In other words, we can reduce this amount by at most 1.

Without loss of generality we assume that β_1 only intersects H_a in a single vertex. Suppose this vertex is in δ_i and consider performing the folds in ϵ_i . After folding ϵ_i , the subsegment of the terminal part of α identified with the initial part of β_1 either heads counterclockwise from v_0 (which we are not concerned with as this adds to the amount by which γ_1 covers a) or it is contained in the union of δ_i and ϵ_i . Indeed, we can just check locally that when folding two edges e_1 and e_2 in ϵ_i together in H_a that β_1 cannot fold past (in the clockwise direction) the image of their terminal vertices. This involves a few cases depending on the relative positions of β_1 , e_1 and e_2 ; all of which are easily verified.

Similarly, if β_1 only intersects H_a in a vertex of ϵ_i , then we find the the subsegment of the terminal portion of α that is identified with β_1 either heads counterclockwise from v_0 or it is contained in ϵ_i .

Thus when performing the folds in $H_a \rightarrow H'_a$, the initial portion of β_1 is identified at most one copy of H_a . Likewise, the same holds for the terminal portion of β_0 . Therefore the image path γ_1 n -covers a . \square

As a consequence of the claim, we have $\ell_{G'}(a^n) \leq BBT(f_1)$. Since H'_a is immersed in every graph along the folding path between G_1 and

G' , we have $\ell_{G'}(a^n) = \text{Lip}(f_1)\ell_{G_1}(a^n)$, so that

$$\ell_{G_1}(a^n) = \ell_{G'}(a^n)/\text{Lip}(f_1) \leq \text{BBT}(f_1)/\text{Lip}(f_1) \leq 1$$

and so $\ell_{G_1}(a) \leq \frac{1}{n}$. \square

Combining Proposition 5.1 with Proposition 3.5 we get the first of the main results of this paper.

Theorem 5.2. *Suppose $G, G' \in CV_k$ with $d = d_L(G, G')$ such that $\tau_a(G, G') \geq n+2$ for some a . Then there is a geodesic $\alpha: [0, d] \rightarrow CV_k$ such that $\alpha(0) = G$ and $\alpha(d) = G'$ and for some $t \in [0, d]$, we have $\ell_{\alpha(t)}(a) \leq 1/n$. In other words, $\alpha([0, d]) \cap CV_k^{1/n}(a) \neq \emptyset$.*

Additionally, combining Proposition 5.1 with Proposition 4.8 we get the second of the main results of this paper.

Theorem 5.3. *Suppose $\phi \in \text{Out } F_k$ is fully irreducible, with unstable tree T_- and lamination Λ_- such that $\tau_a(T_-, \Lambda_-) \geq n+4$ for some $a \in F_k$. Then given any train-track G , there is an axis \mathcal{L}_ϕ for ϕ that contains G and a graph G_0 such that $\ell_{\tilde{G}_0}(a) \leq 1/n$. In other words, $\mathcal{L}_\phi \cap CV_k^{1/n}(a) \neq \emptyset$.*

6. EXAMPLE

Here we present an application of Theorem 5.3 in which we describe the asymptotic behavior of the translation length in CV_k of certain elements of $\text{Out } F_k$, given as products $\phi_n = \delta_1^n \delta_2^{-n}$ of powers of Dehn twists δ_1, δ_2 . These types of outer automorphisms were considered in [13] and used in [12] to show that there is no homological obstruction to full irreducibility. We briefly recall the setup here; for more details consult either of the references [12, 13].

6.1. Dehn twists. A *cyclic* tree is a Bass–Serre tree associated to a splitting of F_k over \mathbb{Z} , either as an amalgamated free product or as an HNN-extension. To such a tree is an associated (outer) automorphism called a *Dehn twist*. Given $F_k = A *_{\langle c \rangle} B$ we define the Dehn twist automorphism δ_c of F_k by:

$$\begin{aligned} \forall a \in A \quad \delta_c(a) &= a \\ \forall b \in B \quad \delta_c(b) &= bcb^{-1}. \end{aligned}$$

Likewise, given $F_k = A *_{\mathbb{Z}} \langle A, t \mid t^{-1}ct = c' \rangle$ for $c, c' \in A$, we define the Dehn twist δ_c of F_k by:

$$\begin{aligned} \forall a \in A \quad \delta_c(a) &= a \\ \delta_c(t) &= ct. \end{aligned}$$

Two cyclic trees T_1 and T_2 *fill* if their associated Dehn twists δ_1, δ_2 do not have any common invariant conjugacy classes of proper free factors or cyclic subgroups. As mentioned above, if T_1 and T_2 fill, then for large enough n , the element $\delta_1^n \delta_2^{-n}$ is fully irreducible (and hyperbolic) [13, Theorem 5.3].

6.2. Currents. A (*measured geodesic*) *current* on F_k is an F_k -invariant and σ -invariant positive Radon measure on $\partial^2 F_k$ (refer to Section 2.6). Such measures were originally considered by Bonahon [9], see also [29]. Given a tree $T \in CV_k$, there is an identification between ∂F_k and ∂T used to interpret a current as a measure on the set of (bi-infinite) geodesics in T . Given a tight path $\alpha \subset T$, the *two-sided cylinder* $Cyl_T(\alpha)$ is the collection of geodesics that contain α ; such sets determine a basis for $\partial^2 T$, and so in turn for $\partial^2 F_k$. When α is a fundamental domain for the action of $a \in F_k$ on $T^{(a)}$, we will denote $Cyl_T(\alpha)$ by $Cyl_T(a)$. For a current $\nu \in Curr(F_k)$, define $\langle a, \nu \rangle_T = \nu(Cyl_T(a))$. As ν is F_k -invariant, this is well-defined. The current is uniquely defined by the values $\langle a, \nu \rangle_T$. If $c \in F_k$ is not a proper power, then we define the *counting current* η_c of c by:

$$\langle a, \eta_c \rangle_T = \# \text{ of axes of conjugates of } c \text{ in } Cyl_T(a)$$

If $b = c^m$ where c is not a proper power, then $\eta_b = m\eta_c$.

Recall that an element $\phi \in \text{Out } F_k$ is *hyperbolic* if it has no nontrivial periodic conjugacy classes in F_k ; all such elements are necessarily non-geometric, in the sense that they are not induced by a surface homeomorphism. A hyperbolic fully irreducible element of $\text{Out } F_k$ acts on the projectivized space of currents $\mathbb{P}Curr(F_k)$ with North-South dynamics [34]. In particular, there are both *stable* $[\mu_+(\phi)]$ and *unstable* $[\mu_-(\phi)]$ fixed projectivized currents associated to such an element. A similar statement holds for non-hyperbolic fully irreducible elements as well, after restricting to the subspace of $\mathbb{P}Curr(F_k)$ consisting just of those currents in the closure of the set of counting currents of primitive elements in F_k [31].

The *support* $Supp(\nu)$ of a current ν is the closure of the union of all open sets U such that $\nu(U) > 0$. The support of a current is a lamination. The relationship between the stable currents and stable laminations of a fully irreducible element of $\text{Out } F_k$ is given by the proposition below. The result is probably well-known, but to our knowledge, its proof does not appear in the literature. See also [30] for closely related results.

Proposition 6.1. *Suppose $\phi \in \text{Out } F_k$ is fully irreducible with stable and unstable laminations Λ_+, Λ_- and stable and unstable currents μ_+, μ_- . We have $\text{Supp}(\mu_\pm) = \Lambda_\pm$.*

Proof. Let $g : G \rightarrow G$ be a train-track representative of ϕ . Let $a \in F_k$ be a primitive element and $\alpha \subset G$ the reduced loop representing its conjugacy class. Then α is the union of N legal paths in G for some N , so that for all $m \geq 0$, the closed path $g^m(\alpha)$ consists of N segments of leaves of the stable lamination Λ_+ .

The set of cylinders $\text{Cyl}_{\tilde{G}}(\gamma)$, γ a reduced path in \tilde{G} , not containing any leaf of Λ_+ give a cover of the complement of Λ_+ . Choose one such cylinder $\text{Cyl}_{\tilde{G}}(\gamma)$, so that γ is not a subsegment of any leaf of Λ_+ . Then for any $m \geq 0$, the reduced loop $[g^m(\alpha)]$ contains at most N copies of the image of γ in G , and hence $\eta_{\phi^m(a)}(\text{Cyl}_{\tilde{G}}(\gamma)) \leq N$. Recall that, because a was chosen to be primitive, we have the convergence of $[\eta_{\phi^m(a)}] \rightarrow [\mu_+]$. Now for a sequence λ_m to give the convergence of $\frac{1}{\lambda_m} \eta_{\phi^m(a)} \rightarrow \mu_+$, it is necessary that $\lambda_m \rightarrow \infty$ [32, Theorem 1.2]. Thus we have $\mu_+(\text{Cyl}_{\tilde{G}}(\gamma)) = 0$.

We have shown that $\text{Supp}(\mu_+) \subseteq \Lambda_+$, a nonempty sublamination of a minimal lamination [7, 32]. The claim of the proposition is verified. \square

6.3. Axes of products of Dehn twists. Let $k \geq 3$ and fix two filling cyclic trees T_1, T_2 with Dehn twists δ_1 and δ_2 . Let c_1, c_2 denote the respective edge stabilizers. We assume that the set $\{c_1, c_2\}$ is not *separable*, i.e., no conjugates of c_1 and c_2 are contained in a proper free factor of F_k , nor in complementary free factors. Further, we assume that c_1 and c_2 are not *simultaneously elliptic* in \overline{CV}_k , i.e., $\ell_T(c_1) + \ell_T(c_2) \neq 0$ for all $T \in \overline{cv}_k$. These conditions can be guaranteed, for instance, by requiring c_1 and c_2 to be primitive elements sufficiently far apart in the free factor complex [14].

For the remainder of this section, elements $\phi_n \in \text{Out } F_k$ denote the outer automorphisms induced by the automorphisms $\delta_1^n \delta_2^{-n}$. For large enough n , the elements ϕ_n are fully irreducible and hyperbolic [13, Theorem 5.3]. From [12, Theorem 5.2], we understand the limiting behavior of the stable and unstable currents: $[\mu_+(\phi_n)] \rightarrow [\eta_{c_1}]$ and $[\mu_-(\phi_n)] \rightarrow [\eta_{c_2}]$. Using this, together with the parabolic behavior of Dehn twists on \overline{CV}_k [15], we show that the sequence of stable and unstable trees likewise converge to the expected trees (cf., [12, Remark 5.3]).

Theorem 6.2. *The trees $T_+(\phi_n) \in \overline{CV}_k$ converge to T_2 . Similarly, the trees $T_-(\phi_n)$ converge to T_1 .*

Proof. Denote $\psi_n = \delta_2^{-n} \delta_1^n$ so that $\phi_n = \delta_2^n \psi_n \delta_2^{-n}$. Then as the outer automorphisms are conjugate by δ_2^n , we have $T_+(\phi_n) = T_+(\psi_n) \delta_2^{-n}$.

Recall that in [12, Theorem 5.2], we determined that $\lim_{n \rightarrow \infty} [\mu_-(\psi_n)] = [\eta_{c_1}]$. The continuity of the Kapovich–Lustig intersection form (see [32] for its definition and properties) implies that c_1 has a fixed point in an accumulation point of the sequence $\{T_+(\psi_n)\}$ (see [12, Remark 5.3]). Therefore as c_1 and c_2 are not simultaneously elliptic, c_2 has positive translation length in any such accumulation point.

As \overline{CV}_k is compact, some subsequence of $\{T_+(\phi_n)\}$ converges. Consider such a convergent subsequence $\{T_+(\phi_{n_m})\} \subseteq \{T_+(\phi_n)\}$. By passing to a further subsequence, we can assume that both $\{T_+(\phi_{n_{m_\ell}})\}$ and $\{T_+(\psi_{n_{m_\ell}})\}$ converge. Let T_∞ denote the limit of the latter sequence. By the above remark, c_2 has positive translation length on the tree T_∞ .

Let $U \subset \overline{CV}_k$ be a neighborhood of T_2 . As the set $\{T_+(\psi_{n_{m_\ell}})\} \cup \{T_\infty\}$ is compact, and as c_2 has positive translation length on every tree therein, by [15, Theorem 13.2], there is an N such that for $\ell \geq N$ we have $T_+(\phi_{n_{m_\ell}}) = T_+(\psi_{n_{m_\ell}}) \delta_2^{-n_{m_\ell}} \in U$. Therefore the subsequence $\{T_+(\phi_{n_m})\}$ converges to T_2 . As this is true for every convergent subsequence of $\{T_+(\phi_n)\}$, and as \overline{CV}_k is compact, we have the convergence of $T_+(\phi_n) \rightarrow T_2$. Applying the same argument to $\phi_n^{-1} = \delta_2^n \delta_1^{-n}$ we see that $T_-(\phi_n) \rightarrow T_1$ as well. \square

Fix bases \mathcal{T}_1 and \mathcal{T}_2 for F_k , obtained from the vertex group(s) (and possibly a choice of stable letter in the case of an HNN-extension) of the Bass-Serre trees T_1 and T_2 , respectively; to see how this is done, we refer to Section 3.1 of [13]. Let $T_{\mathcal{T}_1}$ and $T_{\mathcal{T}_2}$ be the Cayley trees for the basis \mathcal{T}_1 and \mathcal{T}_2 , respectively. See [12, 13] for the details underlying these constructions.

Proposition 6.3. *For sufficiently large n , we have:*

$$\tau_{c_2}(T_{\mathcal{T}_2}, \Lambda_-(\phi_n)) \geq \frac{n}{2}.$$

Proof. The proposition follows from a slight modification of the arguments from Theorem 5.2 in [12]. In its proof (equation (5.9)), we showed that for every $\epsilon > 0$ and integer $r > 0$, there is an $N > 0$ such that for $n \geq N$:

$$\frac{\langle c_2^r, \mu_-(\phi_n) \rangle_{T_{\mathcal{T}_2}}}{\omega_{T_{\mathcal{T}_2}}(\mu_-(\phi_n))} > 1 - \epsilon. \quad (6.1)$$

The $\omega_{T_{\mathcal{T}_2}}(\cdot)$ in the demoninator is a normalization factor whose only relevant value to the present discussion is $\omega_{T_{\mathcal{T}_2}}(\mu_-(\phi_n))$; it may thus

be treated as a positive constant.² Equation (6.1) shows that there is a leaf of $\Lambda_-(\phi_n) = \text{Supp}(\mu_-(\phi_n))$ contained in the cylinder $\text{Cyl}_T(c_2^r)$, and hence $\tau_{c_2}(T_{\mathcal{T}_2}, \Lambda_-(\phi_n)) \geq r$.

Following the same analysis as in [12, Theorem 5.2], fixing $r = n/2$, one can show:

$$\left| 1 - \frac{\langle c_2^{n/2}, \mu_-(\phi_n) \rangle_{T_{\mathcal{T}_2}}}{\omega_{T_{\mathcal{T}_2}}(\mu_-(\phi_n))} \right| < \left| \frac{\frac{1}{2}An^2 + A_1n + A_2}{An^2 - B_1n - B_2} \right| + \frac{\epsilon}{2} \quad (6.2)$$

for some fixed positive constants A, A_1, A_2, B_1 and B_2 .³ Thus for large enough n , we have that $\langle c_2^{n/2}, \mu_-(\phi_n) \rangle_T > 0$, and hence there is a leaf of $\Lambda_-(\phi_n) = \text{Supp}(\mu_-(\phi_n))$ contained in the cylinder $\text{Cyl}_T(c_2^{n/2})$. This implies that $\tau_{c_2}(T_{\mathcal{T}_2}, \Lambda_-(\phi_n)) \geq n/2$, as claimed. \square

We can use the fact that the sequence of trees $\{T_-(\phi_n)\}$ converges to T_1 to show that the twist of $T_-(\phi_n)$ with $\Lambda_-(\phi_n)$ relative to c_2 is also approximately at least n .

Proposition 6.4. *There exists a constant $D \geq 1$ such that for sufficiently large n :*

$$\tau_{c_2}(T_-(\phi_n), \Lambda_-(\phi_n)) \geq \frac{n}{D}.$$

Proof. As before, let $T_{\mathcal{T}_2}$ be the Cayley tree corresponding to the basis \mathcal{T}_2 . By Proposition 6.3, for each n there is a leaf $\ell_n: \mathbb{R} \rightarrow T_{\mathcal{T}_2}$ of $\Lambda_-(\phi_n)$ that intersects the axis of c_2 in a segment of length at least $n\ell_T(c_2)/2$. We must verify that this overlap is not significantly reduced when mapping to $T_-(\phi_n)$.

Fix an F_k -equivariant map $f: T_{\mathcal{T}_2} \rightarrow T_1$ and scale the metric on T_1 so that $\text{Lip}(f) \leq 1$, and thus $\text{BBT}(f) \leq 1$ (we assume that the volume of $T_{\mathcal{T}_2}/F_k$ is 1). By scaling the metrics on $T_-(\phi_n)$ we have the convergence of $T_-(\phi_n) \rightarrow T_1$ from Theorem 6.2. Thus for large enough n , we can choose equivariant maps $f_n: T_{\mathcal{T}_2} \rightarrow T_-(\phi_n)$ so that $\text{Lip}(f_n) \leq 2$ and so $\text{BBT}(f_n) \leq 2$. As convergence is in the space of length functions, and as $\ell_{T_1}(c_2) > 0$, there is $\delta > 0$ such that $0 < \delta < \ell_{T_-(\phi_n)}(c_2) < 1/\delta$ for all n .

Now let $x_n \in \ell_n(\mathbb{R}) \cap T_{\mathcal{T}_2}^{(c_2)}$ be such that $y_n = c_2^{n/2}x_n \in \ell_n(\mathbb{R}) \cap T_{\mathcal{T}_2}^{(c_2)}$. Thus the path $[f_n(x_n), f_n(y_n)]$ contains an arc of the axis of c_2 in $T_-(\phi_n)$ of length at least $\frac{n}{2}\ell_{T_-(\phi_n)}(c_2)$. Further notice that the distance from

²On the other hand, to recognize (6.1) from equation (5.9) in the proof of [12, Theorem 5.2], it should be observed that $\omega_{T_{\mathcal{T}_2}}(\eta_{c_2}) = \langle c_2^r, \eta_{c_2} \rangle_{T_{\mathcal{T}_2}}$.

³Compare this to equation (5.9) from [12, Theorem 5.2] where the numerator of the righthand side is linear in n , and note that the constants β_1, β_2 there depend on r . Here in (6.2), the numerator is quadratic because of the choice of $r = n/2$.

either $f_n(x_n)$ or $f_n(y_n)$ to this arc is at most 2 (an upper bound for the bounded back tracking constant).

As $\ell_n: \mathbb{R} \rightarrow T_{\mathcal{T}_2}$ is a leaf of the unstable lamination, after tightening its image in $T_-(\phi_n)$ we obtain a geodesic $[f_n(\ell_n(\mathbb{R}))]$, and the same statement in the previous paragraph for the segment $[f_n(x_n), f_n(y_n)]$ and the axis of c_2 holds in turn for $[f_n(x_n), f_n(y_n)]$ and the geodesic $[f_n(\ell_n(\mathbb{R}))]$. Hence the leaf of $\Lambda_-(\phi_n)$ whose image in $T_-(\phi_n)$ is $[f_n(\ell_n(\mathbb{R}))]$ intersects the axis of c_2 along a segment of length at least:

$$\begin{aligned} \frac{n\ell_{T_-(\phi_n)}(c_2)}{2} - 4 &> \frac{n\ell_{T_-(\phi_n)}(c_2)}{2} - \frac{4\ell_{T_-(\phi_n)}(c_2)}{\delta} \\ &= \ell_{T_-(\phi_n)}(c_2) \left(\frac{n\delta - 8}{4\delta} \right) \\ &= \ell_{T_-(\phi_n)}(c_2) \frac{n}{D} \end{aligned}$$

for some constant $D > 0$, provided $n > 8/\delta$. Thus

$$\tau_{c_2}(T_-(\phi_n), \Lambda_-(\phi_n)) \geq \frac{n}{D}.$$

□

It follows from the proposition, together with Theorem 5.3, that for each element ϕ_n , there is a train-track map $g_n: G_n \rightarrow G_n$ such that $\ell_{G_n}(c_2) \leq D'/n$ for some constant D' . Note that we are using the fact that every graph on the axis of ϕ represents a train-track of ϕ .

Recall that we assumed that $\{c_1, c_2\}$ is not separable in F_k .

Lemma 6.5. *If $\{c_1, c_2\}$ is not separable, then for large enough n , neither is $\{c_2, \delta_1^n(c_2)\}$.*

Proof. This is easy to see using Whitehead graphs. Since the set $\{c_1, c_2\}$ is not separable, the union of their Whitehead graphs is connected and does not have a cut vertex (in an appropriate basis) [39]. As cancellation is bounded, for large enough n , the subword representing c_1 will appear as a subword of $\delta_1^n(c_2)$. Hence the union of the Whitehead graphs of c_2 and $\delta_1^n(c_2)$ will cover the union of the Whitehead graphs of c_1 and c_2 . In particular, their union will be connected and will not have a cut vertex. This implies the set $\{c_2, \delta_1^n(c_2)\}$ is not separable. □

Note that $\phi_n(c_2) = \delta_1^n(c_2)$ so that, as a consequence of the lemma, every edge in the track-track graph G_n must be crossed by either c_2 or $\phi_n(c_2)$. Therefore, the length of $\phi_n(c_2)$ is at least $1 - D'/n = (n - D')/n$, and thus the Lipschitz constant for ϕ_n is at least

$$\frac{(n - D')/n}{D'/n} = n/D' - 1.$$

In particular, we have now shown that for some constant $K_1 > 0$:

$$\frac{1}{K_1} \log n \leq tr_{CV_k}(\phi_n)$$

where $tr_{CV_k}(\phi) = \min\{d_L(G, G\phi) \mid G \in CV_k\}$ is the minimal translation length of the element ϕ .

We obtain the corresponding upper bound on $tr_{CV_k}(\phi_n)$ by explicitly constructing a path by piecing together geodesic segments such as those constructed in Example 2.2.

As before, let $T_{\mathcal{T}_1}$ and $T_{\mathcal{T}_2}$ be the Cayley trees for the basis \mathcal{T}_1 and \mathcal{T}_2 , respectively. We consider these trees as points in CV_k , with every edge of each tree having length $1/k$. We first connect $T_{\mathcal{T}_2}\delta_2^n$ to $T_{\mathcal{T}_2}$ by a geodesic of length $\sim \log n$. Then we follow an optimal path P from $T_{\mathcal{T}_2}$ to $T_{\mathcal{T}_1}$, and then connect $T_{\mathcal{T}_1}$ to $T_{\mathcal{T}_1}\delta_1^n$ with a geodesic which has length $\sim \log n$. Finally, using the δ_1^n -translate $P\delta_1^n$ of P , we connect $T_{\mathcal{T}_1}\delta_1^n$ to $T_{\mathcal{T}_2}\delta_1^n$ (see Figure 2). As the length of P is independent of n , translating the entire path by δ_2^{-n} , we have for all n :

$$d_L(T_{\mathcal{T}_2}, T_{\mathcal{T}_2}\phi_n) \leq K_2 \log n$$

for some $K_2 > 0$.




FIGURE 2. A path from $T_{\mathcal{T}_2}\delta_2^n$ to $T_{\mathcal{T}_2}\delta_1^n$.

Combining this upper bound with the previous lower bound, we have established the following:

Theorem 6.6. *Let T_1, T_2 be two cyclic trees that fill with associated Dehn twist automorphisms δ_1 and δ_2 and let c_1, c_2 denote the respective edge stabilizers. Suppose that $\{c_1, c_2\}$ is not separable and that c_1 and c_2 are not simultaneously elliptic in $\overline{CV_k}$. For $n \geq 1$, let ϕ_n be the outer automorphism induced by $\delta_1^n \delta_2^{-n}$. Then there is a constant $K = K(T_1, T_2)$ such that for large enough n :*

- (1) *there is a train-track representative $g_n: G_n \rightarrow G_n$ such that $\ell_{G_n}(c_2) \leq K/n$, and*
- (2) *$\frac{1}{K} \log n \leq tr_{CV_k}(\phi_n) \leq K \log n$.*

REFERENCES

- [1] Y. ALGOM-KFIR, *Strongly contracting geodesics in Outer space.* arXiv:math/0812.1555.

- [2] Y. ALGOM-KFIR AND M. BESTVINA, *Asymmetry of Outer Space*. arXiv:math/0910.5408.
- [3] E. ALIBEGOVIĆ, *Translation lengths in $\text{Out}(F_n)$* , *Geom. Dedicata*, 92 (2002), pp. 87–93. Dedicated to John Stallings on the occasion of his 65th birthday.
- [4] J. BEHRSTOCK, M. BESTVINA, AND M. CLAY, *Growth rate of intersection numbers for free group automorphisms*, *J. Topol.*, 3 (2010), pp. 280–310.
- [5] M. BESTVINA, *A Bers-like proof of the existence of train tracks for free group automorphisms*. arXiv:math/1001.0325.
- [6] M. BESTVINA AND M. FEIGHN, *Outer limits*. preprint (1992) <http://andromeda.rutgers.edu/~feighn/papers/outer.pdf>.
- [7] M. BESTVINA, M. FEIGHN, AND M. HANDEL, *Laminations, trees, and irreducible automorphisms of free groups*, *Geom. Funct. Anal.*, 7 (1997), pp. 215–244.
- [8] M. BESTVINA AND M. HANDEL, *Train tracks and automorphisms of free groups*, *Ann. of Math. (2)*, 135 (1992), pp. 1–51.
- [9] F. BONAHO, *Geodesic currents on negatively curved groups*, in *Arboreal group theory* (Berkeley, CA, 1988), vol. 19 of *Math. Sci. Res. Inst. Publ.*, Springer, New York, 1991, pp. 143–168.
- [10] M. R. BRIDSON AND A. HAEFLIGER, *Metric spaces of non-positive curvature*, vol. 319 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*, Springer-Verlag, Berlin, 1999.
- [11] M. CLAY, *Contractibility of deformation spaces of G -trees*, *Algebr. Geom. Topol.*, 5 (2005), pp. 1481–1503 (electronic).
- [12] M. CLAY AND A. PETTET, *Current twisting and nonsingular matrices*. arXiv:math/0907.1075, to appear in *Commentarii Mathematici Helvetici*.
- [13] ———, *Twisting out fully irreducible automorphisms*, *Geom. Funct. Anal.*, 20 (2010), pp. 657–689.
- [14] M. CLAY AND P. REYNOLDS, *Stabilizers of filling free factors*. in preparation.
- [15] M. M. COHEN AND M. LUSTIG, *Very small group actions on \mathbf{R} -trees and Dehn twist automorphisms*, *Topology*, 34 (1995), pp. 575–617.
- [16] T. COULBOIS, A. HILION, AND M. LUSTIG, *\mathbb{R} -trees and laminations for free groups. I. Algebraic laminations*, *J. Lond. Math. Soc. (2)*, 78 (2008), pp. 723–736.
- [17] ———, *\mathbb{R} -trees and laminations for free groups. II. The dual lamination of an \mathbb{R} -tree*, *J. Lond. Math. Soc. (2)*, 78 (2008), pp. 737–754.
- [18] ———, *\mathbb{R} -trees and laminations for free groups. III. Currents and dual \mathbb{R} -tree metrics*, *J. Lond. Math. Soc. (2)*, 78 (2008), pp. 755–766.
- [19] M. CULLER AND J. W. MORGAN, *Group actions on \mathbf{R} -trees*, *Proc. London Math. Soc. (3)*, 55 (1987), pp. 571–604.
- [20] M. CULLER AND K. VOGTMANN, *Moduli of graphs and automorphisms of free groups*, *Invent. Math.*, 84 (1986), pp. 91–119.
- [21] B. FARB, A. LUBOTZKY, AND Y. MINSKY, *Rank-1 phenomena for mapping class groups*, *Duke Math. J.*, 106 (2001), pp. 581–597.
- [22] S. FRANCAVIGLIA AND A. MARTINO, *The isometry group of outer space*. arXiv:0912.0299.
- [23] ———, *Metric properties of outer space*, *Publ. Mat.*, 55 (2011), pp. 433–473.
- [24] V. GUIRADEL, *Cœur et nombre d'intersection pour les actions de groupes sur les arbres*, *Ann. Sci. École Norm. Sup. (4)*, 38 (2005), pp. 847–888.

- [25] V. GUIRARDEL AND G. LEVITT, *The outer space of a free product*, Proc. Lond. Math. Soc. (3), 94 (2007), pp. 695–714.
- [26] U. HAMENSTADT, *Lines of minima in outer space*. arXiv:math/0911.3620.
- [27] M. HANDEL AND L. MOSHER, *The expansion factors of an outer automorphism and its inverse*, Trans. Amer. Math. Soc., 359 (2007), pp. 3185–3208 (electronic).
- [28] ———, *Axes in outer space*, Mem. Amer. Math. Soc., 213 (2011), pp. vi+104.
- [29] I. KAPOVICH, *Currents on free groups*, in Topological and asymptotic aspects of group theory, vol. 394 of Contemp. Math., Amer. Math. Soc., Providence, RI, 2006, pp. 149–176.
- [30] I. KAPOVICH AND M. LUSTIG, *Invariant laminations for irreducible automorphisms of free groups*. arXiv:math/1104.1265.
- [31] ———, *The actions of $\text{Out}(F_k)$ on the boundary of outer space and on the space of currents: minimal sets and equivariant incompatibility*, Ergodic Theory Dynam. Systems, 27 (2007), pp. 827–847.
- [32] ———, *Intersection form, laminations and currents on free groups*, Geom. Funct. Anal., 19 (2010), pp. 1426–1467.
- [33] G. LEVITT AND M. LUSTIG, *Irreducible automorphisms of F_n have north-south dynamics on compactified outer space*, J. Inst. Math. Jussieu, 2 (2003), pp. 59–72.
- [34] R. MARTIN, *Non-Uniquely Ergodic Foliations of Thin Type, Measured Currents and Automorphisms of Free Groups*, PhD thesis, UCLA, 1995.
- [35] H. A. MASUR AND Y. N. MINSKY, *Geometry of the complex of curves. II. Hierarchical structure*, Geom. Funct. Anal., 10 (2000), pp. 902–974.
- [36] Y. MINSKY, *Quasi-projections in teichüller space*, J. Reine Angew. Math., 473 (1996), pp. 121–136.
- [37] K. RAFI, *A characterization of short curves of a Teichmüller geodesic*, Geom. Topol., 9 (2005), pp. 179–202 (electronic).
- [38] ———, *A combinatorial model for the Teichmüller metric*, Geom. Funct. Anal., 17 (2007), pp. 936–959.
- [39] J. R. STALLINGS, *Whitehead graphs on handlebodies*, in Geometric group theory down under (Canberra, 1996), de Gruyter, Berlin, 1999, pp. 317–330.

DEPT. OF MATHEMATICS, ALLEGHENY COLLEGE, MEADVILLE, PA 16335
E-mail address: mclay@alleggheny.edu

DEPT. OF MATHEMATICS, UNIVERSITY OF BRITISH COLUMBIA, VANCOUVER,
 BC V6T 1Z2
E-mail address: alexandra@math.ubc.ca